arXiv:1209.1166v3 [math.DS] 11 Sep 2012

EXAMPLES OF LAGRANGIAN SYSTEM WHOSE α FUNCTION IS NOT DIFFERENTIABLE AT THE BOUNDARY OF THE FLAT

JIANLU ZHANG

Abstract. In this paper, we construct a Lagrangian system of n degrees of freedom whose α function has a n-dimensional flat at the lowest energy level. At the boundary of the flat there are 2^{n-1} points $c_i(i=1,2,\cdots,2^{n-1})$ of which the Mather set $\mathcal{M}(c_i)$ supports at $n-1$ new minimizing measures except the one supported at the minimal fixed point. Besides, at every point c_i, all of the new minimizing measures have non-zero rotation vectors different from each other. Specially, in the case of two degrees of freedom, this example is also stable under small perturbations of potential functions. In the same way we can show that similar phenomenon still happen at an arbitrary energy level.

1. Introduction

In 1991, Mather raised a framework of variational method to deal with Hamiltonian systems and defined α-function and β-function for the first time [Mat]. Since then, many mathematicians have done a lot of research about these two functions and got many deep and precise results, such as [Man],[Mas],[Car],[B-C]. Thanks to their works we now have a clearer understanding of global behavior in general Hamiltonian systems.

Recently, Zheng Yong proved the existence of infinite many minimal homoclinics under a special premise in his paper [Zh]. But we still can not explain the exact dynamical meaning of this premise. We find his premise is deeply related to the property of α-function at the boundary of flat. In fact, the purpose of his premise is to make sure the existence of new measure at the boundary of flat. Our construction proves the existence of Hamiltonian system with new measure coming out at the boundary of flat in arbitrary degrees of freedom case. Moreover, in the case of two degrees of freedom, Hamiltonian systems with this property is stable under small perturbations of potential functions.

In an arbitrary degrees of freedom case we can construct the following example:

$$L(x,\dot{x})=\langle\dot{x},\dot{x}\rangle_g+u_1(x_n)+u_2(x_n)\sum_{i=1}^{n-1}(1-\cos x_i), \tag{1.1}$$

here the precise form of the Riemannian metric g and potential function will be given in the later section. With the help of this example we get our main result as following:

Key words and phrases. Aubry Mather theory, α function, mechanical systems.

Theorem 1.1. *In the case of n degrees of freedom, there exist u_1,u_2 and Riemannian metric g such that the α function of (1.1) has a flat $\mathbb{F}$ of full dimension at the lowest energy level. There are 2^{n-1} points $c_i (i = 1, 2, \cdots, 2^{n-1})$ at the boundary of the flat with respect to which the Mather set $\mathcal{M}(c_i)$ supports at $n-1$ new minimizing measures. Besides, all of the new minimizing measures have non-vanishing rotation vectors different from each other.*

Remark 1.2. By modifying the Remannian metric g we can make all the new minimizing measures ergodic. Then we actually get $n-1$ ergodic minimizing measures besides the one supported on the fixed point.

Remark 1.3. Besides, we can see that in the following section this phenomenon also happens at a higher energy level flat.

Theorem 1.4. *In the case of two degrees of freedom, this example is also stable under perturbations of potential functions.*

We say a property is **open** in the sense of Mañé if there is an open subset $\mathcal{O} \subset C^\infty(M, \mathbb{R})$ such that the property holds for all the Tonelli Lagrangian $L - u, u \in \mathcal{O}$. Then we can easily get the following conclusion:

Corollary 1.5. *The following property is open in the sense of Mañé of a two degrees of freedom case:*
There is a full-dimensional flat $\mathbb{F}$ corresponding to Mañé critical value at the boundary of which the α function is not differentiable.

The paper is outlined as follows. In Section 2, we state the main background information and tools. In Section 3, we give the construction of the examples. We also give the proof of stability in the case of two degrees of freedom and some remarks in Section 4.

2. BASIC ASSUMPTIONS

2.1. **Tonelli Lagrangian.** Let M be a smooth closed manifold. We consider a Lagrangian $L(x, v) \in C^r(TM, \mathbb{R})$ $(r \geq 2)$, which satisfies:

- **convexity** : the Hessian matrix $\frac{\partial^2 L}{\partial v^2}$ is positive definiteness;
- **superlinearity** : $\lim\limits_{\|v\| \longrightarrow \infty} \frac{L(x,v)}{\|v\|} = \infty$ uniformly with respect to $x \in M$;

Usually we take the manifold M as $\mathbb{T}^n$, and $(x, \dot{x})$ is the local coordinate. It is well known that the Euler-Lagrangian equation is given by:

$$\frac{d}{dt}\frac{\partial L}{\partial \dot{x}}(x, \dot{x}) = \frac{\partial L}{\partial x}(x, \dot{x})$$

For $c \in H^1(\mathbb{T}^n, \mathbb{R})$, we also denote the Tonelli Lagrangian by $L - c$, which also has the previous two properties.

2.2. **Mather Theory.** Mather defined the minimizing measure μ as a probability measure which is supported on the orbits of Euler-Lagrangian equation and satisfies $\int L d\mu = \min_\nu \int L d\nu$. So we can define $\alpha(c)$ and its dual $\beta(h)$ function as:

$$\alpha : H^1(M, \mathbb{R}) \to \mathbb{R}, \quad \alpha(c) = -\min_\mu \int_{TM} L - c d\mu.$$

Here μ ranges from the set of invariant probability measures.

$$\beta : H_1(M,\mathbb{R}) \to \mathbb{R}, \quad \beta(h) = \min_{\rho(\mu)=h} \int_{TM} L d\mu.$$

The two functions are both continuous, convex and superlinear. In fact, they are conjugated with each other in the sense of Legendre transformation [Mat],[R]. From basic convex analysis knowledge we know that if there exists some point on the boundary of the flat at which $\alpha(c)$ is not differentiable, then there must be extra minimizing measures with non vanishing rotation vectors coming out. Hence, we just need to construct an example whose α function is not differentiable at some point of the boundary of the flat.

3. Construction of the example

Use the following example we can get our result in Theorem 1.1.

3.1. **the lowest energy level case.** We construct the Lagrangian as follows.

$$L(x,\dot{x}) = \langle \dot{x},\dot{x}\rangle_g + u_1(x_n) + u_2(x_n)\sum_{i=1}^{n-1}(1-\cos x_i)$$

where $(x,\dot{x}) \in \mathbb{T}^n \times \mathbb{R}^n$ and $x = (x_1, x_2, \ldots, x_n)$. $u_1(x_n)$, $u_2(x_n)$ and the Riemannian metric g are constructed as follows.

First, we make several channels on the $\mathbb{T}^n$ along the n-th coordinate x_n.

$$\begin{cases} \text{channel } A : \mathbb{T}^n \times [-\frac{\pi}{4n}, \frac{\pi}{4n}], \ \mathbb{T} = \mathbb{R}/2\pi, \\ \text{channel } B_i : \mathbb{T}^n \times [\frac{2\pi i}{n} - \frac{\pi}{4n}, \frac{2\pi i}{n} + \frac{\pi}{4n}], \ i = 1, 2, \cdots n-1, \\ \text{channel } C_j : \mathbb{T}^n \times [\frac{(2j-1)\pi}{n} - \frac{\pi}{4n}, \frac{(2j-1)\pi}{n} + \frac{\pi}{4n}], \ j = 1, 2, \cdots n, \end{cases}$$

obviously, channel B_i and C_j are just constructed by shifting channel A along the x_n coordinate. It is easy to see that A and B_i are separated by C_j.

$$u_1(x_n) = \begin{cases} 0, \quad x_n \text{ in channel } A, \\ K \gg 1, \quad x_n \text{ in channel } C_j \text{ and } j = 1, 2, \cdots n, \\ 0 < \delta_i \ll K, \quad x_n \text{ in channel } B_i \text{ and } i = 1, 2, \cdots n-1, \end{cases} \tag{3.1}$$

where the values of δ_i will be given later (in fact it is enough to take $0 < \delta_i \leq \frac{1}{2}$) and $u_1(x_n)$ can be made smooth enough.

$$u_2(x_n) = \begin{cases} 1, \quad x_n \text{ in channel } A, \\ 0, \quad x_n \text{ out of some small neighborhood of channel A}, \end{cases} \tag{3.2}$$

and we can also make $u_2(x_n)$ smooth enough.

At last, we construct the Riemannian metric $\langle \dot{x},\dot{x}\rangle_g = \sum_{i=1}^{n} a_i(x)^2 \dot{x}_i^2$ as follows:

$$G = \begin{pmatrix} a_1^2(x) & 0 & \cdots & 0 \\ 0 & a_2^2(x) & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & a_n^2(x) \end{pmatrix}_{n\times n}$$

is the metric matrix which is diagonal. For x_n in channel A,

$$G = \mathbf{Id}_{n\times n}$$

For x_n in channel C_j, $j = 1, 2, \dots, n$,

$$G = \begin{pmatrix} K & 0 & \cdots & 0 \\ 0 & K & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & K \end{pmatrix}_{n\times n}$$

For x_n in channel B_k, $k = 1, 2, \dots, n-1$,

$$G = \begin{pmatrix} \frac{1}{8} & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \frac{1}{8} & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & 0 & \ddots & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & \frac{1}{8} & 0 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 0 & \frac{1}{4} & 0 & \cdots & 0 \\ 0 & \cdots & \cdots & \cdots & 0 & \frac{1}{8} & 0 & 0 \\ 0 & \cdots & \cdots & \cdots & \cdots & 0 & \ddots & 0 \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & \frac{1}{8} \end{pmatrix}_{n\times n},$$

where $\frac{1}{4}$ is in the k-th row and k-th column.

The projection along the n-th coordinate $p : \mathbb{T}^n \longrightarrow \mathbb{T}^{n-1}$ induces a homomorphism of the cohomology map $p^* : H^1(\mathbb{T}^{n-1}, \mathbb{R}) \longrightarrow H^1(\mathbb{T}^n, \mathbb{R})$. Let $\mathcal{H} = Im(p^*)$ as the subset of the $H^1(\mathbb{T}^n, \mathbb{R})$ with the n-th coordinate $c_n = 0$.

We denote the restriction of Lagrangian L on the channels A, B_i and C_j by L_A, L_{B_i} and L_{C_j}. Also we could use $\alpha_A(c)$, $\alpha_{B_i}(c)$ and $\alpha_{C_j}(c)$ to denote the α function of L restricted to A, B_i, C_j separately, here $c \in \mathcal{H}$.

Moreover, we could calculate $\alpha_A(c)$, $\alpha_{B_i}(c)$ and $\alpha_{C_j}(c)$.

$$\begin{aligned} \alpha_A(c) &= -\min \int L_A - c d\mu, \\ &= -\min \int \langle v, v\rangle_g + \sum_{i=1}^{n-1}(1-\cos x_i) - \sum_{i=1}^{n-1} c_i v_i d\mu, \\ &= -\min \int \sum_{i=1}^{n-1}(v_i^2 + (1-\cos x_i) - c_i v_i) d\mu, \\ &= -\min \sum_{i=1}^{n-1} \int (v_i^2 + (1-\cos x_i) - c_i v_i) d\mu, \\ &= -\sum_{i=1}^{n-1} \min \int (v_i^2 + (1-\cos x_i) - c_i v_i) d\mu. \end{aligned}$$

It is easy to see that L_A consists of $n-1$ independent pendulum systems. Hence, α_A has a flat at $[-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}]^{n-1} \subseteq \mathcal{H}$.

$$\begin{aligned}\alpha_{B_i}(c) &= -\min \int L_{B_i} - c d\mu \\ &= -\min \int \langle v, v\rangle_g + \delta_i - \sum_{i=1}^{n-1} c_i v_i d\mu \\ &= -\delta_i + \sum_{k=1}^{n-1} 16c_k^2 - 12c_i^2, \quad i = 1, 2, \cdots, n-1.\end{aligned}$$

In fact, if we take $\delta_i < \frac{1}{2}$, the level set $\alpha_{B_i}^{-1}(0) \ \subseteq [-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}]^{n-1} \subseteq \mathcal{H}$.

The following lemma gives the relationship between α function of L on $\mathbb{T}^n$ and α function of L restricted to channel A and B_i.

Lemma 3.1.

$$\alpha_L(c) = \max_{i=1,2,\cdots,n-1}\{\alpha_A, \alpha_{B_i}\},$$

where $c = (c_1, c_2, \cdots, c_{n-1}) \in \mathcal{H}$. *This Lemma is still right under small perturbations of potential functions*

Proof. First, we have $\alpha_L(c) \geq \alpha_A(c)$, and $\alpha_L(c) \geq \alpha_{B_i}(c)$. It follows since the action of the local minimizer is bigger than the global minimizer. Hence,

$$\alpha_L(c) \geq \max_{i=1,2,\cdots,n-1}\{\alpha_A, \alpha_{B_i}\}.$$

On the other hand, from [Mn] we know that every minimizing measure can be approximated by a sequence of measures supported on absolutely continuous closed curves, so we just need to certify that every closed curve γ in $\mathbb{T}^n$ has a larger action of the Lagrangian $L - c^*$ than its projection γ^* to $\mathbb{T}^{n-1}$ in some channel A or B_i, here $c^* = (c_1, c_2, \cdots, c_{n-1}, 0) \in \mathcal{H}$. It follows from

$$\begin{aligned}\int_\gamma L - c^* - \int_{\gamma^*} L - c^* &\geq \int a_n^2\dot{\gamma}_n^2 + u_1(x_n) \\ &\geq 0.\end{aligned}$$

Hence,

$$\alpha_L(c) \leq \max_{i=1,2,\cdots,n-1}\{\alpha_A, \alpha_{B_i}\}.$$

If another Lagrangian L' is C^1 close to L, i.e. $L' = L + V(x)$ and $\|V\|_r \leq \epsilon \ll \delta_i$, $i = 1, 2, \cdots, n-1$. Now the channels A, B_i and C_j will be deformed with an magnitude not bigger than ϵ. So these new channels denoted by A', B_i' and C_j' have the same homology group with A, B_i and C_j each. Without loss of generality, we can denote these channels' ϵ neighborhoods by A, B_i and C_j and avoid the deformation.

If the closed curve γ is just in channel A, this Lemma is still right. If the n-th component of the homology type of γ is not zero, it must cross A and at least one

B_i, then we project the part outside of channel A to channel A with the $n-th$ coordinate keeping constant, which is denoted by $(\gamma\cap A^c)^*$ and:

$$\begin{aligned}\int_{\gamma\cap A^c} L'-c^* - \int_{(\gamma\cap A^c)^*} L'-c^* &\geq \int_{\gamma\cap A^c} K_1\dot{\gamma}_n^2+K_2\\ &\geq K_2(T_2-T_1)+\frac{K_1}{T_2-T_1}\int_{T_1}^{T_2}\dot{\gamma}_n^2dt\int_{T_1}^{T_2}1dt\\ &\geq K_2(T_2-T_1)+\frac{K_1}{T_2-T_1}(\int_{T_1}^{T_2}\dot{\gamma}_ndt)^2\\ &\geq 2\sqrt{K_1K_2}\cdot width(C_i)\\ &> 0,\end{aligned}$$

here K_1 and K_2 could be both positive and sufficient large. In fact, $K_1=K^2$ and $K_2=K-\epsilon\gg\frac{K}{2}$ if we take ϵ sufficiently small. The width of some channel C_i is here because A and B_i are separated by at least one C_i. We denote the two endpoints of $(\gamma\cap A^c)^*$ by a, b. Obviously these two points are both in channel A. Then we connect these two points with a geodesic curve ζ and $(\gamma\cap A^c)^*\cup\zeta\cup(\gamma\cap A)\doteq\xi$ form a closed curve. But we know $dist(a,b)\leq\sqrt{n}$, and

$$\begin{aligned}\int_0^1 L'(\zeta,\dot{\zeta})-\langle c^*,\dot{\zeta}\rangle dt &\leq \epsilon+1+\int_0^1|\dot{\zeta}|^2dt\\ &\leq n+1+\epsilon.\end{aligned}$$

Here we could always choose a proper ζ satisfying $\int_0^1\langle c^*,\dot{\zeta}\rangle\geq 0$ because $\mathbb{T}^n$ is a closed manifold. Then we could take sufficient large K such that $n+1+\epsilon\leq 2\sqrt{K_1K_2}\cdot width(C_i)$. So ξ has smaller action value than γ, that means there exists a measure in channel A with smaller action value. Then γ must have a homology type with zero n-th component.

With the help of this, we can directly project $\gamma\cap A^c$ to channel A with the $n-th$ coordinate keeping constant and we needn't extra curve to make it closed. Obviously this curve in A has a smaller action value.So if μ is the minimizing measure of L at C^* and γ is the orbit in its support, we must have either $\gamma\subset A$ or $\gamma\subset A^c$.

For the other case $\gamma\subset A^c$, we can give a similar proof and project γ into some channel B_i with n-th coordinate constant. Since all the possible cases are finite many, we can always take a sufficient large K. Therefore, the lemma is proved. □

Remark 3.2. The stability property under perturbation in this lemma will be used in the next section. Also we can see [S] for more details about this Lemma.

Now, we have constructed our first example of n degrees of freedom. First, $\alpha_L^{-1}(0)$ is a lowest flat of $n-1$ dimensions, here $c\in\mathcal{H}$. This is because $\alpha_{B_i}(c)=-\delta_i+\sum_{k=1}^{n-1}16c_k^2-12c_i^2$ and $\alpha_{B_i}^{-1}(0)\subseteq[-\frac{4\sqrt{2}}{\pi},\frac{4\sqrt{2}}{\pi}]^{n-1}\subseteq\mathcal{H}$.

Second, we could find 2^{n-2} pairs of points which are diametrical with each other at the boundary of the flat. Also at these points $\alpha_L(c)$ is not differentiable along at least $n-1$ different directions, since $\alpha_{B_i}(c)$ cross the flat $[-\frac{4\sqrt{2}}{\pi},\frac{4\sqrt{2}}{\pi}]^{n-1}$ from inner and $\alpha_L(c)=\max_{i=1,2,\cdots,n-1}\{\alpha_A,\alpha_{B_i}\}$. More precisely, the coordinates of these points in $\mathcal{H}$ are easy to write down when ($\delta_i \equiv \frac{1}{2}$):

$$\underbrace{(\pm\frac{1}{\sqrt{8(4n-7)}}, \pm\frac{1}{\sqrt{8(4n-7)}}, \cdots, \pm\frac{1}{\sqrt{8(4n-7)}})}_{n-1}.$$

At each point, there are $n-1$ more measures besides the fix points in channel A. Notice that there is not only one fixed point in the channel A, but we can shrink channel A to just one line $\{x_n = 0\}$ since x_n ia a free variable. Also we can see that the extra $n-1$ new measures are all Lagrangian invariant torus consist of periodic orbits, but we can change the Riemannian metric in the channel B_i to make all the new measures quasi-periodic KAM torus.

At last, we show that the lowest level flat is indeed of full dimension. That's because in the interior of the flat, we have:

$$\widetilde{\mathcal{M}}(c) = \widetilde{\mathcal{N}}(c), \quad \forall c \in int\mathbb{F},$$

and we also knows that $\widetilde{\mathcal{M}}(c)$ just consists of the fixed points. So we could make use of the upper semi-continuous property of the $\widetilde{\mathcal{N}}(c)$ and expand the flat along the directions of $H_1(M, \mathcal{A}(0), \mathbb{R})$. We also recall that all the sublevel set $\{c|\alpha(c) \leq h\}$ is a convex set. So in fact we get a full dimensional flat at the lowest level. See [Zh] for more details, but for the sake of completeness we also give our proof below:

Lemma 3.3. *If $c \in \mathcal{H}^{\perp} \subseteq H_1(\mathbb{T}^n, \mathbb{R})$, there exists a sufficient small $\lambda \ll 1$, such that there exists a flat $[-\lambda c, \lambda c]$ along the direction of c.*

Proof. By the upper semi-continuity of set-valued function $c \longrightarrow \widetilde{\mathcal{N}}(c)$, we know for sufficiently small λ, $\mathcal{N}(\lambda c)$ is still in channel A because $\mathcal{M}(0) = \mathcal{A}(0) = \mathcal{N}(0) = \{0\} \in \mathbb{R}^n$. Then:

$$-\alpha(\lambda c) = \int L - \lambda c d\mu_{\lambda c} = \int L d\mu_{\lambda c} - \langle \lambda c, [\mu_{\lambda c}]\rangle = \int L d\mu_{\lambda c} \geq \int L d\mu_0 = -\alpha(0),$$

so we have $\alpha(\pm\lambda c) \leq \alpha(0)$, where the $-\lambda c$ case is the same with λc. We get $\alpha(\pm\lambda c) = \alpha(0)$ because of the convexity, and then $\alpha_L(c)$ has a full dimensional flat at the lowest energy level. ☐

3.2. **the higher energy level case.** Now, we show that the same phenomenon can happen at a higher energy level.

For the sake of simplicity, we set $n = 3$ and it's easy to generalize the example to a higher dimensional case($n \geq 3$). We could take the Lagrangian as:

$$L(x, \dot{x}) = \frac{1}{2}a_1(x)(\dot{x}_1 - 1)^2 + \frac{1}{2}a_2(x)\dot{x}_2^2 + \frac{1}{2}a_3(x)\dot{x}_3^2 + u_1(x_3) + u_2(x_3)(1 - cosx_2)$$

We just need set two C-type channels , one A-type channel and one B-type channel. The settings of u_1 and u_2 is the same with the previous subsection. Since we still restrict the cohomology to $\mathcal{H}$, so we have $\dot{x}_3 \equiv 0$ and:

$$L_{C_i} \geq \frac{1}{2}(\dot{x}_1 - 1)^2 + K, \quad i = 1, 2,$$

$$L_B = \frac{1}{2}(\dot{x}_1 - 1)^2 + \dot{x}_2^2 + \delta,$$

and

$$L_A = \frac{1}{2}(\dot{x}_1 - 1)^2 + \frac{1}{2}\dot{x}_2^2 + (1 - \cos x_2).$$

Then Lemma 3.1 is still valid:

$$\alpha_L(c)\big|_{c_3=0} = \max\{\alpha_A, \alpha_B\}.$$

Based on our calculation and $c_3 = 0$, the set $\{\alpha_A = 0\}$ is $(0, [-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}], 0)$, which is a flat of one dimension. And $\{\alpha_B = 0\}$ is an elliptical curve with expression as

$$\frac{1}{2}(c_1 + 1)^2 + \frac{1}{4}c_2^2 = \frac{1}{2} + \delta.$$

Restrict to $c_1 = 0$ and we find that the elliptical curve go across $\{\alpha_A = 0\}$ from inner with the intersection points' coordinates $(0, \pm 2\sqrt{\delta}, 0)$. For the purpose of showing that α_L is not differentiable at this two points (in fact we just need show that for one point), we need to calculate the directional derivative along $\overrightarrow{e}_2$.

$$\partial^+_{\vec{e}_2}\alpha_L(0, 2\sqrt{\delta}, 0) = \partial_{\vec{e}_2}\alpha_B(0, 2\sqrt{\delta}, 0) = \sqrt{\delta},$$

and on the other hand

$$\partial^-_{\vec{e}_2}\alpha_L(0, 2\sqrt{\delta}, 0) = \partial_{\vec{e}_2}\alpha_A(0, 2\sqrt{\delta}, 0) = 0.$$

So $\alpha_L(0, 2\sqrt{\delta}, 0)$ is not differentiable along the direction $\vec{e}_2$. That means at this point $(0, 2\sqrt{\delta}, 0)$, there exists an extra minimizing measure besides the closed curves $\mathbb{T} \times \{0\} \times \{0\}$ which is of a different homology.

Remark 3.4. With the upper semi-continuous property of the $\widetilde{\mathcal{N}}(c)$, we can see that in the case of three degrees of freedom the flat is in fact of two dimensions[Zh]. The following graph shows the exact situation of the flat.



Figure 1.

4. Stability of the two degrees of freedom case

In this section we prove the conclusion of Theorem 1.4 in the case of two degrees of freedom.

We take $\epsilon \ll \tau^2 \ll \delta \ll 1$, and shrink the width of channel A and B to the order of $O(\epsilon)$ for the sake of simplity, and τ is used to control the value of potential function. Later we will explain their usage precisely. A ball in $C^r(M,\mathbb{R})$ centered at "0" with a radius of ϵ is denoted by $B(0,\epsilon)$. Any perturbation function $V(x)$ in it can't break the intersection property of α_A and α_B but just deform them of a ϵ order change (see the following graphs). This is because the continuity of α-function with respect to Lagrangian L.



Figure 2.

So in a small neighborhood of point a, there is still an intersection point a' of $\alpha_{L',A}$ and $\alpha_{L',B}$, here $L' = L + V$. With the help of Legendre transformation, the Hamiltonian H' dual to L' is still a quadratic mechanical system, so the α function is axial symmetry. So we just deal with a' point and we can also get the same conclusion of b'. As is known, the maximal potential value is equal to $\alpha'_{L',A}(0)$, which is the minimizing value of $\alpha'_{L',A}$ function[Car]. Without lose of generality, we suppose this value is still 0. If not, we can add a constant to $L + V$ without influencing the property of α function.

Still restrict to $\mathcal{H}$, the curve $\widehat{a'o}$ may be no longer a straight line, we will consider the two different cases separately in the following and disprove both of them. From Lemma 3.1, we know $\widetilde{\mathcal{M}}(a')$ have at least two ergodic minimizing measures, one in channel A and the other in channel B. We denote the one in channel A by $\mu_{a',A}$ and the other $\mu_{a',B}$. From the construction of channel A, the support of this measure can only be a periodic orbit with a rotation vector $(h_1, 0)$ because of the restriction of homology.

If $h_1 = 0$, we could see that the curve $\widehat{a'o}$ is in fact a straight line and from [Mas] we know $\widetilde{\mathcal{A}}(0) \subseteq \widetilde{\mathcal{A}}(a')$. But $\epsilon \ll \delta$, so $\mu_{a',B}$ couldn't have a trivial homology because of the convexity of the $\alpha_{L',B}$. So we get that not only $\widehat{a'ob'}$ is straight, but also $\alpha_{L'}$ is not differential along $\mathcal{H}$ direction. So we get the stability of the example.

If $h_1 > 0$(the same with $h_1 < 0$), we let $a' \longrightarrow a''$, here a'' is the first point with $\alpha_{L',A}(a'') = 0$. Here the case $a'' = 0$ is possible. Then we know that $[a'', 0]$ is a flat and $\widetilde{\mathcal{M}}(0) \subseteq \widetilde{\mathcal{A}}(a'')$, and $\widetilde{\mathcal{A}}(a'') \setminus \widetilde{\mathcal{M}}(0) \neq \emptyset$ [Zh]. Then there must exist a homoclinic orbit or a periodic orbit in $\widetilde{\mathcal{A}}(a'')$.

For the homoclinic orbit case, because the maximal points of potential value are contained in a neighborhood of $(0, 0)$ of a radius of order $O(\epsilon)$ (that's why we



FIGURE 3.

shrink the width of channel A,B), the minimizing homoclinic is asymptotic to this neighborhood $B(0,\epsilon)$. For any $c \in [a', a'']$, $\widetilde{\mathcal{M}}(c)$ can only be made up of periodic



FIGURE 4.

orbits in channel A. Under the weak topology of probability measures, there exists a sequence $c_n \longrightarrow a''$ and $\mu_{L',c_n} \rightharpoonup \mu_{L',0} \in \widetilde{\mathcal{M}}(0)$. And the minimizing homoclinic orbit γ is contained in the Hausdorff limit of $supp\{\mu_{c_n}\}$. As γ is a static orbit in $\widetilde{\mathcal{A}}(a'')$, so

$$\int_{-\infty}^{+\infty} L'(\gamma,\dot\gamma)dt = \langle a'', [\gamma]\rangle, \tag{4.1}$$

here the $\langle , \rangle$ is the inner product induced by de Rham. In fact, $\langle a'', [\gamma]\rangle$ is really a scalar product because the homology of channel A and $a'' \in \mathcal{H}$. As $\epsilon \ll \tau^2 \ll \delta \ll 1$, we have

$$\begin{aligned}\int_{-\infty}^{+\infty} L'(\gamma,\dot\gamma)dt =& \int_{-\infty}^{+\infty} \langle \dot\gamma,\dot\gamma\rangle_g + u_1(\gamma_2(t)) \\ &+ u_2(\gamma_2(t))(1-\cos(\gamma_1(t))) + V(\gamma(t))dt \\ \geq& \int_{-T}^{+T} (\dot\gamma_1{}^2 + \dot\gamma_2{}^2) + u_1(\gamma_2(t)) \\ &+ u_2(\gamma_2(t))(1-\cos(\gamma_1(t))) + V(\gamma(t))dt,\end{aligned} \tag{4.2}$$

here $\gamma\mid_{[-T,T]}$ is the part of γ outside the $O(\tau)$ neighborhood of $(0,0)$. Recall that $\epsilon \ll \tau^2 \ll \delta \ll 1$, we can get a positive lower bound of potential function out of $O(\tau)$ neighborhood of $(0,0)$. As we have supposed $\max_{x\in A} -u_1(x) - u_2(x)(1-$

$\cos(x)) - V(x) = 0$, and the width of channel A is of order $O(\epsilon)$, we have:

$$\begin{aligned}\int_{-T}^{+T} (\dot{\gamma_1}^2 + \dot{\gamma_2}^2) + &u_1(\gamma_2(t)) + u_2(\gamma_2(t))(1-\cos(\gamma_1(t))) + V(\gamma(t))dt \\ &\geq \int_{-T}^{+T} \dot{\gamma_1}^2 + 1 - \cos\tau - \epsilon dt \\ &= \int_{\tau}^{2\pi-\tau} \dot{\gamma}_1 + \frac{1-\cos\tau-\epsilon}{\dot{\gamma}_1} d\gamma_1 \\ &\geq 2(2\pi - 2\tau)\sqrt{1-\cos\tau-\epsilon},\end{aligned}$$

here (γ_1, γ_2) is the coordinate of γ.

At the same time, we have:

$$\begin{aligned}\langle a'', [\gamma]\rangle &= a''\pi_1([\gamma]) \\ &= a'' \\ &\geq 2(2\pi - 2\tau)\sqrt{1-\cos\tau-\epsilon},\end{aligned}$$

because $[\gamma] \equiv (1,0)$. We already have $a'' < a'$. Specially we take $\tau = \frac{1}{8}\pi$, then $2(2\pi - 2\tau)\sqrt{1-\cos\tau-\epsilon} > \frac{\pi}{9}$. But the Riemannian metric can be modified in channel B to make $a' < \frac{\pi}{9}$ and there will be a contradiction to our assumption.

For the periodic orbit case, we can give a same proof as above. And we just need to modify the integral lower and upper bounds to finite in (4.1) and (4.2) to get a same contradiction. So we get the stability property of the example in the case of two degrees of freedom.

Acknowledgement The author thanks Prof. C.Q.Cheng and W.Cheng for talking about the details, and thanks L.Wang for suggesting this problem to me.

References

[B-C] Bernard P. Contreras G. *A generic property of families of Lagrangian systems.* Annals of Math. 167(2008) 1099-1108.

[Car] M.J.Dias Carneiro. *On minimizing measures of the action of autonomous Lagrangians.* Nonlinearity, 8(6):1077-1085,1995.

[Man] R.Mañé. *Generic properties and problems of minimizing measures of Lagrangian systems.* Nonlinearity. **9**(1996), 273-310.

[Mas] Daniel Massart *On Aubry sets and Mather's action functional.* Israel Journal of Mathematics 134(2003),157-171.

[Mat] J.N.Mather. *Action minimizing invariant measures for postive definite Lagrangian systems.* Math. Z. **207** (1991), 169-207.

[R] R.T.Rockafellar. *Convex Analysis.* Princeton University Press, 1970.

[S] Radu Saghin. *On the number of ergodic minimizing measures for Lagrangian flows.* Discrete and Continuous Dynamical Systems. Volume 17, Number 3, March 2007, 501-507.

[Zh] Yong Zheng. *Homoclinic orbits of positive definite Lagrangian systems.* J.Differential Equations 229(2006) , 297-316.

Department of Mathematics, Nanjing University, Nanjing, China, 210093

E-mail address: `jellychung1987@gmail.com`

arXiv:1209.1166v3 [math.DS] 11 Sep 2012

EXAMPLES OF LAGRANGIAN SYSTEM WHOSE α FUNCTION IS NOT DIFFERENTIABLE AT THE BOUNDARY OF THE FLAT

JIANLU ZHANG

Abstract. In this paper, we construct a Lagrangian system of n degrees of freedom whose α function has a lowest n-dimensional flat. At the boundary of the flat there are 2^{n-1} points $c_i(i = 1, 2, \cdots, 2^{n-1})$ of which the Mather set $\mathcal{M}(c_i)$ supports at $n-1$ new minimizing measures except the one supported at the minimal fixed point. Besides, at every point c_i, all of the new minimizing measures have non-zero rotation vectors different from each other. Specially, in the case of two degrees of freedom, this example is also stable under small perturbations of potential functions. In the same way we can show that similar phenomenon still happen at an arbitrary energy level.

1. Introduction

In 1991, Mather raised a framework of variational method to deal with Hamiltonian systems and defined α-function and β-function for the first time [Mat]. Since then, many mathematicians have done a lot of research about these two functions and got many deep and precise results, such as [Man],[Mas],[Car],[B-C]. Thanks to their works we now have a clearer understanding of global behavior in general Hamiltonian systems.

Recently, Zheng Yong proved the existence of infinite many minimal homoclinics under a special premise in his paper [Zh]. But the rationality of this premise is still not well explained. We find his premise is deeply related to the property of α-function at the boundary of flat. In fact, the purpose of his premise is to make sure the existence of new measure at the boundary of flat. Our construction proves the existence of Hamiltonian system with new measure coming out at the boundary of flat in arbitrary degrees of freedom case. Moreover, in the case of two degrees of freedom, Hamiltonian systems with this property is stable under small perturbations of potential functions.

In an arbitrary degrees of freedom case we can construct the following example:

$$L(x, \dot{x}) = \langle \dot{x}, \dot{x} \rangle_g + u_1(x_n) + u_2(x_n) \sum_{i=1}^{n-1} (1 - \cos x_i), \tag{1.1}$$

1991 *Mathematics Subject Classification.* Primary 37Jxx; Secondary 70Hxx.

Key words and phrases. Aubry Mather theory, α function, mechanical systems.

This work was partially supported by the National Basic Research Program of China (Grant No. 2007CB814800) and Natural Scientific Foundation of China (Grant No. 10971093).

here the precise form of the Riemannian metric g and potential function will be given in the later section. With the help of this example we get our main result as following:

Theorem 1.1. *In the case of n degrees of freedom, there exist u_1,u_2 and Riemannian metric g such that the α function of (1.1) has a flat $\mathbb{F}$ of full dimension at the lowest energy level. There are 2^{n-1} points $c_i (i = 1, 2, \cdots , 2^{n-1})$ at the boundary of the flat with respect to which the Mather set $\mathcal{M}(c_i)$ supports at $n-1$ new minimizing measures. Besides, all of the new minimizing measures have non-vanishing rotation vectors different from each other.*

Remark 1.2. By modifying the Remannian metric g we can make all the new minimizing measures ergodic. Then we actually get $n-1$ ergodic minimizing measures besides the one supported on fixed point.

Remark 1.3. Besides, we can see that in the following section this phenomenon also happens at a higher energy level flat.

Theorem 1.4. *In the case of two degrees of freedom, this example is also stable under perturbations of potential functions.*

We say a property is **open** in the sense of Mañé if there is a open subset $\mathcal{O} \subset C^\infty(M, \mathbb{R})$ such that the property holds for all the Tonelli Lagrangian $L-u, u \in \mathcal{O}$. Then we san easily get the following conclusion:

Corollary 1.5. *The following property is open in the sense of Mañé of a two degrees of freedom case:*
There is a full-dimensional flat $\mathbb{F}$ corresponding to Mañé critical value at the boundary of which the α function is not differentiable.

The paper is outlined as follows. In Section 2, we state the main background information and tools. In Section 3, we give the construction of the examples. We also give the proof of stability in the case of two degrees of freedom and some remarks in Section 4.

2. BASIC ASSUMPTIONS

2.1. **Tonelli Lagrangian.** Let M be a smooth closed manifold. We consider a Lagrangian $L(x, v) \in C^r(TM, \mathbb{R})$ $(r \geq 2)$, which satisfies:

- **convexity** : the Hessian matrix $\frac{\partial^2 L}{\partial v^2}$ is positive definiteness;
- **superlinearity** : $\lim\limits_{\|v\| \longrightarrow \infty} \frac{L(x,v)}{\|v\|} = \infty$ uniformly with respect to $x \in M$;

Usually we take the manifold M as $\mathbb{T}^n$, and $(x, \dot{x})$ is the local coordinate. It is well known that the Euler-Lagrangian equation is given by:

$$\frac{d}{dt}\frac{\partial L}{\partial \dot{x}}(x, \dot{x}) = \frac{\partial L}{\partial x}(x, \dot{x})$$

For $c \in H^1(\mathbb{T}^n, \mathbb{R})$, we also denote the Tonelli Lagrangian by $L - c$, which also has the previous three properties.

2.2. **Mather Theory.** Mather defined the minimizing measure μ as a probability measure which is supported on the orbits of Euler-Lagrangian equation and satisfies $\int L d\mu = \min_\nu \int L d\nu$. So we can denote the α function $\alpha(c)$ and its dual $\beta(h)$ function as:

$$\alpha : H^1(M, \mathbb{R}) \to \mathbb{R}, \quad \alpha(c) = -\min_{\mu} \int_{TM} L - c d\mu.$$

Here μ is a invariant probability measure.

$$\beta : H_1(M, \mathbb{R}) \to \mathbb{R}, \quad \beta(h) = \min_{\rho(\mu)=h} \int_{TM} L d\mu.$$

The two functions are both continuous, convex and superlinear. In fact, they are conjugated with each other in the sense of Legendre transformation [Mat],[R]. From basic convex analysis knowledge we know that if there exists some point on the boundary of the flat at which the $\alpha(c)$ function is not differentiable, then there must be extra minimizing measures with non vanishing rotation vectors coming out. Hence, we just need to construct an example whose α function is not differentiable at some point of the boundary of the flat.

3. Construction of the example

3.1. **the lowest energy level case.** We construct the Lagrangian as follows.

$$L(x, \dot{x}) = \langle \dot{x}, \dot{x} \rangle_g + u_1(x_n) + u_2(x_n) \sum_{i=1}^{n-1} (1 - \cos x_i)$$

where $(x, \dot{x}) \in \mathbb{T}^n \times \mathbb{R}^n$ and $x = (x_1, x_2, \dots, x_n)$. $u_2(x_n)$, $u_1(x_n)$ and the Riemannian metric g are constructed as follows.

First, we make several channels on the $\mathbb{T}^n$ along the n-th coordinate x_n.

$$\begin{cases} \text{channel } A : \mathbb{T}^n \times [-\frac{\pi}{4n}, \frac{\pi}{4n}], \ \mathbb{T} = \mathbb{R}/2\pi, \\ \text{channel } B_i : \mathbb{T}^n \times [\frac{2\pi i}{n} - \frac{\pi}{4n}, \frac{2\pi i}{n} + \frac{\pi}{4n}], \ i = 1, 2, \cdots n-1, \\ \text{channel } C_j : \mathbb{T}^n \times [\frac{(2j-1)\pi}{n} - \frac{\pi}{4n}, \frac{(2j-1)\pi}{n} + \frac{\pi}{4n}], \ j = 1, 2, \cdots n, \end{cases}$$

obviously, channel B_i and C_j are just constructed by shifting channel A along the x_n coordinate. It is easy to see that A and B_i are separated by C_j.

$$u_1(x_n) = \begin{cases} 0, \quad x_n \text{ in channel } A, \\ K \gg 1, \quad x_n \text{ in channel } C_j \text{ and } j = 1, 2, \cdots n, \\ 0 < \delta_i \ll K, \quad x_n \text{ in channel } B_i \text{ and } i = 1, 2, \cdots n-1, \end{cases} \tag{3.1}$$

where the values of δ_i will be given later (in fact it is enough to take $0 < \delta \leq \frac{1}{2}$) and $u_1(x_n)$ can be made smooth enough.

$$u_2(x_n) = \begin{cases} 1, \quad x_n \text{ in channel } A, \\ 0, \quad x_n \text{ out of some small neighborhood of channel A}, \end{cases} \tag{3.2}$$

and we can also make $u_2(x_n)$ smooth enough.

At last, we construct the Riemannian metric $\langle \dot{x}, \dot{x} \rangle_g = \sum_{i=1}^n a_i(x)^2 \dot{x}_i^2$ as follows:

$$G = \begin{pmatrix} a_1^2(x) & 0 & \cdots & 0 \\ 0 & a_2^2(x) & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & a_n^2(x) \end{pmatrix}_{n\times n}$$

is the metric matrix which is diagonal. For x_n in channel A,

$$G = \mathbf{Id}_{n\times n}$$

For x_n in channel C_j, $j = 1, 2, \dots, n$,

$$G = \begin{pmatrix} K & 0 & \cdots & 0 \\ 0 & K & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & K \end{pmatrix}_{n\times n}$$

For x_n in channel B_k, $k = 1, 2, \dots, n-1$,

$$G = \begin{pmatrix} \frac{1}{8} & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & \frac{1}{8} & 0 & \cdots & \cdots & 0 \\ 0 & 0 & \frac{1}{4} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & \frac{1}{8} & 0 & 0 \\ 0 & \cdots & \cdots & 0 & \ddots & 0 \\ 0 & \cdots & \cdots & \cdots & 0 & \frac{1}{8} \end{pmatrix}_{n\times n},$$

where $\frac{1}{4}$ is on the k-th row and k-th column.

The projection along the n-th coordinate $p : \mathbb{T}^n \longrightarrow \mathbb{T}^{n-1}$ induces a homomorphism of the cohomology map $p^* : H^1(\mathbb{T}^{n-1}, \mathbb{R}) \longrightarrow H^1(\mathbb{T}^n, \mathbb{R})$. Let $\mathcal{H} = Im(p^*)$ as the subset of the $H^1(\mathbb{T}^n, \mathbb{R})$ with the n-th coordinate $c_n = 0$.

We denote the restriction of Lagrangian L on the channels A, B_i and C_j by L_A,L_{B_i} and L_{C_j}. Also we could use $\alpha_A(c)$, $\alpha_{B_i}(c)$ and $\alpha_{C_j}(c)$ to denote the α function of L restricted to A, B_i, C_j separately, here $c \in \mathcal{H}$.

Moreover, we could calculate $\alpha_A(c)$, $\alpha_{B_i}(c)$ and $\alpha_{C_j}(c)$.

$$\begin{aligned} \alpha_A(c) &= -\min \int L_A - c d\mu, \\ &= -\min \int \langle v, v \rangle_g + \sum_{i=1}^{n-1} (1 - \cos x_i) - \sum_{i=1}^{n-1} c_i v_i d\mu, \\ &= -\min \int \sum_{i=1}^{n-1} (v_i^2 + (1 - \cos x_i) - c_i v_i) d\mu, \\ &= -\min \sum_{i=1}^{n-1} \int (v_i^2 + (1 - \cos x_i) - c_i v_i) d\mu, \\ &= -\sum_{i=1}^{n-1} \min \int (v_i^2 + (1 - \cos x_i) - c_i v_i) d\mu. \end{aligned}$$

It is easy to see that L_A consists of $n-1$ independent pendulum systems. Hence, α_A has a flat at $[-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}]^{n-1} \subseteq \mathcal{H}$.

$$\begin{aligned}\alpha_{B_i}(c) &= -\min \int L_{B_i} - c d\mu \\ &= -\min \int \langle v, v\rangle_g + \delta_i - \sum_{i=1}^{n-1} c_i v_i d\mu \\ &= -\delta_i + \sum_{k=1}^{n-1} 16c_k^2 - 12c_i^2, \quad i = 1, 2, \cdots, n-1.\end{aligned}$$

In fact, if we take $\delta_i < \frac{1}{2}$, the level set $\alpha_{B_i}^{-1}(0) \subseteq [-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}]^{n-1} \subseteq \mathcal{H}$.

The following lemma gives the relationship between α function of L on $\mathbb{T}^n$ and α function of L restricted to channel A and B_i.

Lemma 3.1.

$$\alpha_L(c) = \max_{i=1,2,\cdots,n-1} \{\alpha_A, \alpha_{B_i}\},$$

where $c = (c_1, c_2, \cdots, c_{n-1}) \in \mathcal{H}$. This Lemma is still right under small perturbations of potential functions

Proof. First, we have $\alpha_L(c) \geq \alpha_A(c)$, and $\alpha_L(c) \geq \alpha_{B_i}(c)$. It follows since the action of the local minimizer is bigger than the global minimizer. Hence,

$$\alpha_L(c) \geq \max_{i=1,2,\cdots,n-1} \{\alpha_A, \alpha_{B_i}\}.$$

On the other hand, from [Mn] we know that every minimizing measure can be approximated by a sequence of measures supported on absolutely continuous closed curves, so we just need to certify that every closed curve γ in $\mathbb{T}^n$ has a larger action of the Lagrangian $L - c^*$ than its projection γ^* to $\mathbb{T}^{n-1}$ in some channel A or B_i, here $c^* = (c_1, c_2, \cdots, c_{n-1}, 0) \in \mathcal{H}$. It follows from

$$\begin{aligned}\int_\gamma L - c^* - \int_{\gamma^*} L - c^* &\geq \int a_n^2 \dot{\gamma}_n^2 + u_1(x_n) \\ &\geq 0.\end{aligned}$$

Hence,

$$\alpha_L(c) \leq \max_{i=1,2,\cdots,n-1} \{\alpha_A, \alpha_{B_i}\}. \qquad (*)$$

If another Lagrangian L' is C^1 close to L, i.e. $L' = L + V(x)$ and $\|V\|_r \leq \epsilon \ll \delta_i$,$i = 1, 2, \cdots, n-1$. If the closed curve γ is just in channel A, $(*)$ is still right. If γ crosses completely one B_i, then we project the part outside of channel A to channel A with the $n-th$ coordinate constant, which is denoted by $(\gamma\cap A^c)^*$ and:

$$\begin{aligned}\int_{\gamma\cap A^c} L' - c^* - \int_{(\gamma\cap A^c)^*} L' - c^* &\geq \int_{\gamma\cap A^c} C_1\dot{\gamma}_n^2 + C_2\\ &\geq C_2(T_2-T_1) + \frac{C_1}{T_2-T_1}\int_{T_1}^{T_2}\dot{\gamma}_n^2 dt\int_{T_1}^{T_2} 1dt\\ &\geq C_2(T_2-T_1) + \frac{C_1}{T_2-T_1}(\int_{T_1}^{T_2}\dot{\gamma}_n dt)^2\\ &\geq 2\sqrt{C_1C_2}\cdot width(B_i)\\ &\geq 0,\end{aligned}$$

here C_1 and C_2 could be both positive if we take ϵ sufficiently small. Therefore, the lemma is proved. □

Remark 3.2. The stability property under perturbation in this lemma will be used in the next section. Also we can see [S] for more details about this Lemma.

Now, we have constructed our first example of n degrees of freedom. First, $\alpha_L^{-1}(0)$ is a lowest flat of $n-1$ dimensions, here $c\in\mathcal{H}$. This is because $\alpha_{B_i}(c) = -\delta_i + \sum_{k=1}^{n-1} 16c_k^2 - 12c_i^2$ and $\alpha_{B_i}^{-1}(0)\subseteq[-\frac{4\sqrt{2}}{\pi},\frac{4\sqrt{2}}{\pi}]^{n-1}\subseteq\mathcal{H}$.

Second, we could find 2^{n-2} pairs of points which are diametrical with each other at the boundary of the flat. Also at these points the $\alpha_L(c)$ funtion is not differentiable along at least $n-1$ different directions, since the $\alpha_{B_i}(c)$ functions cross the flat $[-\frac{4\sqrt{2}}{\pi},\frac{4\sqrt{2}}{\pi}]^{n-1}$ downward and $\alpha_L(c) = \max_{i=1,2,\cdots,n-1}\{\alpha_A,\alpha_{B_i}\}$. The following graph shows the α function on the section of $\mathcal{H}$:

More precisely, the coordinates of these points are easy to write when ($\delta_i\equiv\frac{1}{2}$):

$$\underbrace{(\pm\frac{1}{\sqrt{8(4n-7)}},\pm\frac{1}{\sqrt{8(4n-7)}},\cdots,\pm\frac{1}{\sqrt{8(4n-7)}})}_{n-1}.$$

At each point, there are $n-1$ more measures besides the fix points in channel A. Notice that there is not only one fixed point in the channel A, but we can shrink channel A to just one line $\{x_n=0\}$ since x_n ia a free variable. Also we can see that the extra $n-1$ new measures are all Lagrangian invariant torus consist of periodic orbits, but we can change the Riemannian metric in the channel B_i to make all the new measures quasi-periodic KAM torus.

At last, we show that the lowest level flat is indeed of full dimensions. That's because in the interior of the flat, we have:

$$\widetilde{\mathcal{M}}(c) = \widetilde{\mathcal{N}}(c), \quad \forall c\in int\mathbb{F},$$

and we also knows that $\widetilde{\mathcal{M}}(c)$ just consists of the fixed points. So we could make use of the upper semi-continuous property of the $\widetilde{\mathcal{N}}(c)$ and expand the flat along the directions of $H_1(M,\mathcal{A}(0),\mathbb{R})$. We also recall that all the level set $\{c|\alpha(c)\leq h\}$ is a convex set. So in fact we get a full dimensional flat at the lowest level. See [Zh] for more details, but for the sake of completeness we also give our proof below:

Lemma 3.3. *If $c\in\mathcal{H}^\perp\subseteq H_1(\mathbb{T}^n,\mathbb{R})$, there exists sufficient small $\lambda\ll 1$, such that there exists a flat $[-\lambda c,\lambda c]$ along the direction of c.*

Proof. By the upper semi-continuity of set-valued function $c \longrightarrow \widetilde{\mathcal{N}}(c)$, we know for sufficiently small λ, $\mathcal{N}(\lambda c)$ is still in channel A because $\mathcal{M}(0) = \mathcal{A}(0) = \mathcal{N}(0) = \{0\} \in \mathbb{R}^n$. Then:

$$-\alpha(\lambda c) = \int L - \lambda c d\mu_{\lambda c} = \int L d\mu_{\lambda c} - \langle \lambda c, [\mu_{\lambda c}] \rangle = \int L d\mu_{\lambda c} \geq \int L d\mu_0 = -\alpha(0),$$

so we have $\alpha(\pm\lambda c) \leq \alpha(0)$, where the $-\lambda c$ case is the same with λc. We get $\alpha(\pm\lambda c) = \alpha(0)$ because of the convexity, and then $\alpha_L(c)$ has a full dimensional flat at the lowest energy level. ☐

3.2. **the higher energy level case.** Now, we show that the same phenomenon can happen at a higher energy level.

For the sake of simplicity, we set $n = 3$ and it's easy to generalize the example to a higher dimensional case($n \geq 3$). We could take the Lagrangian as:

$$L(x, \dot{x}) = \frac{1}{2} a_1(x)(\dot{x}_1 - 1)^2 + \frac{1}{2} a_2(x)\dot{x}_2^2 + \frac{1}{2} a_3(x)\dot{x}_3^2 + u_1(x_3) + u_2(x_3)(1 - cos x_2)$$

We just need set two C-type channels , one A-type channel and one B-type channel. The settlement of u_1 and u_2 is the same with the previous subsection. Since we still restrict the cohomology to $\mathcal{H}$, so we have $\dot{x}_3 \equiv 0$ and:

$$L_{C_i} \geq \frac{1}{2}(\dot{x}_1 - 1)^2 + K, \quad i = 1, 2,$$

$$L_B = \frac{1}{2}(\dot{x}_1 - 1)^2 + \dot{x}_2^2 + \delta,$$

and

$$L_A = \frac{1}{2}(\dot{x}_1 - 1)^2 + \frac{1}{2}\dot{x}_2^2 + (1 - \cos x_2).$$

Then Lemma 3.1 is still valid:

$$\alpha_L(c)\big|_{c_3=0} = \max\{\alpha_A, \alpha_B\}.$$

Based on our calculation and the restriction $c_3 = 0$, the set $\{\alpha_A = 0\}$ is $(0, [-\frac{4\sqrt{2}}{\pi}, \frac{4\sqrt{2}}{\pi}], 0)$, which is a flat of one dimension. And $\{\alpha_B = 0\}$ is an elliptical curve with expression as

$$\frac{1}{2}(c_1 + 1)^2 + \frac{1}{4}c_2^2 = \frac{1}{2} + \delta.$$

Restrict to $c_1 = 0$ and we find that the elliptical curve go across $\{\alpha_A = 0\}$ from inner with the intersection points' coordinates $(0, \pm 2\sqrt{\delta}, 0)$. For the purpose of showing that α_L is not differentiable at this two points (in fact we just need show that for one point), we need to calculate the directional derivative along $\overrightarrow{e}_2$.

$$\partial^+_{\vec{e}_2} \alpha_L(0, 2\sqrt{\delta}, 0) = \partial_{\vec{e}_2} \alpha_B(0, 2\sqrt{\delta}, 0) = \sqrt{\delta},$$

and on the other hand

$$\partial^-_{\vec{e}_2} \alpha_L(0, 2\sqrt{\delta}, 0) = \partial_{\vec{e}_2} \alpha_A(0, 2\sqrt{\delta}, 0) = 0.$$

So $\alpha_L(0, 2\sqrt{\delta}, 0)$ is not differentiable along the direction $\vec{e}_2$. That means at this point $(0, 2\sqrt{\delta}, 0)$, there exists an extra minimizing measure besides the closed curves $\mathbb{T} \times \{0\} \times \{0\}$ which is of a different homology.

Remark 3.4. With the upper semi-continuous property of the $\widetilde{\mathcal{N}}(c)$, we can see that in the case of three degrees of freedom the flat is in fact of two dimensions[Zh]. The following graph shows the exact situation of the flat.



Figure 1.

4. Stability of the two degrees of freedom case

In this section we prove the stability of the example under perturbation of potential functions in the case of two degrees of freedom. We take $\epsilon \ll \tau^2 \ll \delta \ll 1$, and shrink the width of channel A and B to the order of $O(\epsilon)$ for the sake of simplity, and τ is used to control the potential function's value. Later we will explain their usage precisely. A ball in $C^r(M, \mathbb{R})$ centered at "0" with a radius of ϵ is denoted by $B(0, \epsilon)$. Any perturbation function $V(x)$ in it can't break the intersection property of α_A and α_B but just deform them of a ϵ order change. This is because the continuity of α-function with respect to Lagrangian L (see the following graphs).



Figure 2.

So in a small neighborhood of point a, there is still an intersection point a' of $\alpha_{L',A}$ and $\alpha_{L',B}$, here $L' = L + V$. With the help of Legendre transformation, the Hamiltonian H' dual to L' is still a quadratic mechanical system, so the α function is axial symmetry. So we just deal with a' point and we can also get the same conclusion of b'. As is known, the maximal potential value is equal to $\alpha'_{L',A}(0)$, which is the minimizing value of $\alpha'_{L',A}$ function[Car]. Without lose of generality, we suppose this value is still 0. If not, we can add a constant to $L + V$ without influencing the property of α function.

Still restrict to $\mathcal{H}$, the curve $\widehat{a'o}$ may be no longer a straight line, we will consider the straight case and $\alpha_{L',A}(a') > \alpha_{L',A}(0) = 0$ case separately in the following and both disprove them. From Lemma 3.1, we know $\widetilde{\mathcal{M}}(a')$ have at least two ergodic minimizing measures, one in channel A and the other in channel B. We denote the one in channel A by $\mu_{a',A}$ and the other $\mu_{a',B}$. From the construction of channel A, the support of this measure can only be a periodic orbit with a rotation vector $(h_1, 0)$ because of the restriction of homology.

If $h_1 = 0$, we could see that the curve $\widehat{a'o}$ is in fact a straight line and from [Mas] we know $\widetilde{\mathcal{A}}(0) \subseteq \widetilde{\mathcal{A}}(a')$. But $\epsilon \ll \delta$, so $\mu_{a',B}$ couldn't have a trivial homology because of the convexity of the $\alpha_{L',B}$. So we get that not only $\widehat{a'ob'}$ is straight, but also $\alpha_{L'}$ is not differential along $\mathcal{H}$ direction. So we get the stability of the example.

If $h_1 > 0$(the same with $h_1 < 0$), we let $a' \longrightarrow a''$, here a'' is the first point with $\alpha_{L',A}(a'') = 0$. Here the case $a'' = 0$ is possible. Then we know that $[a'', 0]$ is a flat and $\widetilde{\mathcal{M}}(0) \subseteq \widetilde{\mathcal{A}}(a'')$, and $\widetilde{\mathcal{A}}(a'') \setminus \widetilde{\mathcal{M}}(0) \neq \emptyset$[Zh]. Then there must exist a homoclinic orbit or a periodic orbit in $\widetilde{\mathcal{A}}(a'')$.



FIGURE 3.

For the homoclinic case, because the maximal value points contain in a neighborhood of $(0, 0)$ of a radius of order $O(\epsilon)$ (that's why we shrink the width of channel A,B), the minimizing homoclinic is asymptotic to this neighborhood $B(0, \epsilon)$. For



FIGURE 4.

any $c \in [a', a'']$, $\widetilde{\mathcal{M}}(c)$ can only be made up of periodic orbits in channel A. Under the weak topology of probability measures, there exists a sequence $c_n \longrightarrow a''$ and $\mu_{L',c_n} \rightharpoonup \mu_{L',0} \in \widetilde{\mathcal{M}}(0)$. And the minimizing homoclinic orbit γ is contained in the Hausdorff limit of $supp\{\mu_{c_n}\}$. As γ is a static orbit in $\widetilde{\mathcal{A}}(a'')$, so

$$\int_{-\infty}^{+\infty} L'(\gamma, \dot{\gamma})dt = \langle a'', [\gamma] \rangle, \tag{4.1}$$

here the $\langle , \rangle$ is the inner product induced by de Rham. In fact, $\langle a'', [\gamma] \rangle$ is really a scalar product because the homology of channel A and $a'' \in \mathcal{H}$. As $\epsilon \ll \tau^2 \ll \delta \ll 1$, we have

$$\begin{aligned}\int_{-\infty}^{+\infty} L'(\gamma,\dot\gamma)dt =&\int_{-\infty}^{+\infty}\langle\dot\gamma,\dot\gamma\rangle_g + u_1(\gamma_2(t))\\ &+u_2(\gamma_2(t))(1-\cos(\gamma_1(t)))+V(\gamma(t))dt\\ \geq&\int_{-T}^{+T}({\dot\gamma_1}^2+{\dot\gamma_2}^2)+u_1(\gamma_2(t))\\ &+u_2(\gamma_2(t))(1-\cos(\gamma_1(t)))+V(\gamma(t))dt,\end{aligned} \tag{4.2}$$

here $\gamma\mid_{[-T,T]}$ is the part of γ outside the $O(\tau)$ neighborhood of $(0,0)$. Recall that $\epsilon \ll \tau^2 \ll \delta \ll 1$, we can get a positive lower bound of potential function out of $O(\tau)$ neighborhood of $(0,0)$. As we have supposed $\max_{x\in A} -u_1(x) - u_2(x)(1-\cos(x)) - V(x) = 0$, and the width of channel A is of order $O(\epsilon)$, we have:

$$\begin{aligned}\int_{-T}^{+T}({\dot\gamma_1}^2+{\dot\gamma_2}^2)&+u_1(\gamma_2(t))+u_2(\gamma_2(t))(1-\cos(\gamma_1(t)))+V(\gamma(t))dt\\ &\geq\int_{-T}^{+T}{\dot\gamma_1}^2+1-\cos\tau-\epsilon dt\\ &=\int_{\tau}^{2\pi-\tau}\dot\gamma_1+\frac{1-\cos\tau-\epsilon}{\dot\gamma_1}d\gamma_1\\ &\geq 2(2\pi-2\tau)\sqrt{1-\cos\tau-\epsilon},\end{aligned}$$

here (γ_1,γ_2) is the coordinate of γ.

At the same time, we have:

$$\begin{aligned}\langle a'',[\gamma]\rangle =&a''\pi_1([\gamma])\\ =&a''\\ \geq&2(2\pi-2\tau)\sqrt{1-\cos\tau-\epsilon},\end{aligned}$$

because $[\gamma]\equiv(1,0)$. We already have $a''<a'$. Specially we take $\tau=\frac{1}{8}\pi$, then $2(2\pi-2\tau)\sqrt{1-\cos\tau-\epsilon} > \frac{\pi}{9}$. But the Riemannian metric can be modified in channel B to make $a'<\frac{\pi}{9}$ and there will be a contradiction to our assumption.

For the periodic orbit case, we can give a same proof as above. And we just need to modify the integral lower and upper bounds to finite in (4.1) and (4.2) to get a same contradiction. So we get the stability property of the example in the case of two degrees of freedom.

Acknowledgement The author thanks Prof. C.Q.Cheng and W.Cheng for talking about the details and suggestions.

References

[B-C] Bernard P. Contreras G. *A generic property of families of Lagrangian systems.* Annals of Math. 167(2008) 1099-1108.

[Car] M.J.Dias Carneiro. *On minimizing measures of the action of autonomous Lagrangians.* Nonlinearity, 8(6):1077-1085,1995.

[Man] R.Mañé. *Generic properties and problems of minimizing measures of Lagrangian systems.* Nonlinearity. **9**(1996), 273-310.

[Mas] Daniel Massart *On Aubry sets and Mather's action functional.* Israel Journal of Mathematics 134(2003),157-171.

- [Mat] J.N.Mather. *Action minimizing invariant measures for postive definite Lagrangian systems*. Math. Z. **207** (1991), 169-207.
- [R] R.T.Rockafellar. *Convex Analysis*. Princeton University Press, 1970.
- [S] Radu Saghin. *On the number of ergodic minimizing measures for Lagrangian flows*. Discrete and Continuous Dynamical Systems. Volume 17, Number 3, March 2007, 501-507.
- [Zh] Yong Zheng. *Homoclinic orbits of positive definite Lagrangian systems*. J.Differential Equations 229(2006) , 297-316.

Department of Mathematics, Nanjing University, Nanjing, China, 210093

E-mail address: `jellychung1987@gmail.com`